

**ehp**

**ENVIRONMENTAL  
HEALTH  
PERSPECTIVES**

ehponline.org

**Global Farm Animal Production and  
Global Warming: Impacting and  
Mitigating Climate Change**

---

**Gowri Koneswaran and Danielle Nierenberg**

**doi:10.1289/ehp.11034 (available at <http://dx.doi.org/>)  
Online 31 January 2008**



**NIEHS**  
National Institute of  
Environmental Health Sciences

National Institutes of Health  
U.S. Department of Health and Human Services

## Global Farm Animal Production and Global Warming: Impacting and Mitigating Climate Change

Gowri Koneswaran (1) and Danielle Nierenberg (1, 2)

(1) The Humane Society of the United States, Washington, D.C.

(2) Worldwatch Institute, Washington, D.C.

### **Send Proofs to:**

Gowri Koneswaran

HSUS—Farm Animal Welfare

2100 L St., N.W.

Washington, DC 20037

Tel: 202.676.2362

Fax: 202.676.2372

[gkoneswaran@humanesociety.org](mailto:gkoneswaran@humanesociety.org)

## **Running Title: Global Farm Animal Production and Climate Change**

**Key Words:** animal agriculture, climate change, concentrated animal feeding operation (CAFO), diet, environment, farm animals, farm animal welfare, food choices, global warming, greenhouse gas emissions (GHGs)

### **Acknowledgments:**

Both authors are staff members of The Humane Society of the United States. Danielle Nierenberg also serves as a Senior Fellow with the Worldwatch Institute, where she was previously employed.

### **Abbreviations:**

CH<sub>4</sub>: methane

CO<sub>2</sub>: carbon dioxide

FAO: Food and Agriculture Organization of the United Nations

GHG: greenhouse gas

GWP: global warming potential

IPCC: Intergovernmental Panel on Climate Change

N<sub>2</sub>O: nitrous oxide

NCAR: National Center for Atmospheric Research

USDA: U.S. Department of Agriculture

U.S. EPA: U.S. Environmental Protection Agency

**Outline:**

Abstract

Background

Global Warming and Climate Change

Anthropogenic Influences

Objectives

Discussion

Impacts of Growing Livestock Populations and Intensifying Production

CO<sub>2</sub> Emissions from Animal Agriculture

Nitrogen from Fertilizer and Feed Production

Methane and Nitrous Oxide

Conflict, Hunger, and Disease

Strategies and Next Steps

Conclusion

References

## **Abstract**

**BACKGROUND:** The farm animal sector is the single largest anthropogenic user of land, contributing to many environmental problems, including global warming and climate change.

**OBJECTIVES:** To synthesize and expand upon existing data on this sector's contribution to climate change.

**METHODS:** We analyzed the scientific literature and documented this sector's GHG emissions, as well as various mitigation strategies.

**DISCUSSIONS:** An analysis of meat, egg, and milk production encompasses not only the direct rearing and slaughtering of animals, but also grain and fertilizer production for animal feed, waste storage and disposal, water use, and energy expenditures on farms and in transporting feed and finished animal products, among other key impacts of the production process as a whole.

**CONCLUSIONS:** Immediate and far-reaching changes in current animal agriculture practices and consumption patterns are both critical and timely if GHGs from this sector are to be mitigated.

## **Introduction**

Though much evidence of its negative impacts on environmental integrity, community sustainability, public health, and animal welfare has been amassed, the totality of the animal agriculture sector's global impacts has remained largely underestimated and underappreciated. A recent review of the relevant data calculated the sector's contributions to global greenhouse gas (GHG) emissions and determined them to be so significant that, measured in carbon dioxide equivalent, the animal agriculture sector's emissions surpass those of the transportation sector (Steinfeld et al. 2006).

### ***Global Warming and Climate Change***

The three main GHGs are carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), and nitrous oxide (N<sub>2</sub>O) (Steinfeld et al. 2006). While most attention has focused on carbon dioxide, methane and nitrous oxide—both extremely potent GHGs—have greater global warming potentials (GWPs) than CO<sub>2</sub>. By assigning CO<sub>2</sub> a value of one GWP, the warming potentials of these other gases can be expressed on a CO<sub>2</sub>-equivalent basis (Paustian et al. 2006; Steinfeld et al. 2006): CH<sub>4</sub> has a GWP of 23 while N<sub>2</sub>O has a GWP of 296.

Global warming's many impacts are already detectable. As glaciers retreat, the sea level rises, the tundra thaws, hurricanes and other extreme weather events occur more frequently, and penguins, polar bears, and other species struggle to survive (Topping 2007), experts anticipate even greater increases in the intensity and prevalence of these changes as the 21<sup>st</sup> century brings rises in GHG emissions. The five warmest years since the 1890s were 1998, 2002, 2003, 2004, and 2005 (NASA 2006). Indeed, average global temperatures have risen considerably (IPCC 2007c), and the Intergovernmental Panel on Climate Change (IPCC) predicts increases of 1.8-3.9°C (3.2-7.1°F) by 2100 (IPCC 2007c). The

IPCC's latest report, released in November 2007, warns that climate change "could lead to some impacts that are abrupt or irreversible" (IPCC 2007b).

### ***Anthropogenic Influences***

Although some natural occurrences contribute to GHG emissions (IPCC 2007c), the overwhelming consensus among the world's most reputable climate scientists contends that human activities are culpable for a majority of this increase in temperature (IPCC 2007a). An IPCC report issued in April 2007 concluded "with high confidence that anthropogenic warming over the last three decades has had a discernible influence on many physical and biological systems" (IPCC 2007a).

While transportation and the burning of fossil fuels have typically been regarded as the chief contributors to GHG emissions and climate change, a November 2006 report, *Livestock's Long Shadow: Environmental Issues and Options*, by the Food and Agriculture Organization (FAO) of the United Nations highlighted the farm animal production sector's substantial role. Identifying it as "a major threat to the environment" (FAO 2006), the FAO found that the animal agriculture sector emits 18 percent, or nearly one-fifth, of human-induced GHG emissions, more than cars, SUVs, and other vehicles (Steinfeld et al. 2006).

### **Objectives**

The authors' objective is to outline the animal agriculture sector's share of global GHG emissions by synthesizing and expanding upon the data in *Livestock's Long Shadow* with more recent reports by the IPCC, data from the U.S. Environmental Protection Agency (EPA), and studies on GHGs from agriculture and mitigation strategies (Cederberg and Stadig 2003; IFOAM 2004; IPCC 2007a, 2007b,

2007c; McMichael et al. 2007; Ogino et al. 2007; U.S. EPA 2007a; Verge et al. 2007). We also draw attention to links between this sector and the far-reaching impacts of climate change on conflict, hunger, and disease, while underscoring the roles of animal agriculture industries, policymakers, and individual consumers in mitigating this sector's contributions to climate change and global warming.

## **Discussion**

### ***Impacts of Growing Livestock Populations and Intensifying Production***

According to FAOSTAT, globally, approximately 56 billion land animals are reared and slaughtered for human consumption annually, and livestock inventories are expected to double by 2050, with most increases occurring in the developing world (Steinfeld et al. 2006). As the numbers of farm animals reared for meat, egg, and dairy production rise, so do their GHG emissions. The U.S. Department of Agriculture (USDA 2004) has noted that “GHG emissions from livestock are inherently tied to livestock population sizes because the livestock are either directly or indirectly the source for the emissions.” Since the 1940s, for example, escalating farm animal populations—in large, confined operations, in particular—have significantly increased methane emissions from both animals and their manure (Paustian et al. 2006).

In recent decades, increasing numbers of animals are raised in intensive production systems, in which chickens, pigs, turkeys, and other animals are confined in cages, crates, pens, stalls, and warehouse-like grow-out facilities. These production systems are devoid of environmental stimuli, adequate space or means by which to experience most natural behaviors. Furthermore, because these industrialized, “landless” facilities tend to produce more manure than can be used as fertilizer on nearby cropland

(FAO 2005b), manure is instead “distributed to a small, local landmass resulting in soil accumulation and runoff of phosphorus, nitrogen, and other pollutants” (Thorne 2007).

While extensive, or pasture-based farming methods remain the norm in Africa and some parts of Asia, the trend in Latin America and Asia is to increasingly favor intensive production systems over more sustainable and welfare-friendly practices (Nierenberg 2006). According to a 2007 report describing GHG emissions from agriculture, “[i]n recent years, industrial livestock production has grown at twice the rate of more traditional mixed farming systems and at more than six times the rate of production based on grazing” (Verge et al. 2007).

Confining greater numbers of animals indoors and further separating production operations from agricultural land will only exacerbate the environmental problems already posed by this sector, which the FAO has deemed “one of the top two or three most significant contributors to the most serious environmental problems, at every scale from local to global” (Steinfeld et al. 2006).

### ***CO<sub>2</sub> Emissions from Animal Agriculture***

Regarded as the most important GHG, CO<sub>2</sub> has the most significant direct warming impact on global temperature due to the sheer volume of its emissions. Of all the natural and human-induced influences on climate over the past 250 years, the largest is due to increased CO<sub>2</sub> concentrations attributed to burning fossil fuels and deforestation (Bierbaum et al. 2007).

The animal agriculture sector accounts for approximately 9 percent of total CO<sub>2</sub> emissions (Steinfeld et al. 2006), which are primarily the result of fertilizer production for feed crops, on-farm energy

expenditures, feed transport, animal product processing and transport, and land use changes (Steinfeld et al. 2006).

Burning fossil fuels to produce fertilizers for feed crops may emit 41 million tonnes of CO<sub>2</sub> per year (Steinfeld et al. 2006), with 1 tonne equalling 1 metric ton, or 1,000 kilograms. Vast amounts of artificial nitrogenous fertilizer are used to grow farm animal feed, primarily comprised of corn and soybeans. Most of this fertilizer is produced in factories dependent on fossil-fuel energy (Steinfeld et al. 2006). The Haber-Bosch process, which produces ammonia in order to create nitrogen-based artificial fertilizer, is used to produce 100 million tonnes of fertilizer for feed crops annually (Steinfeld et al. 2006).

An additional 90 million tonnes of CO<sub>2</sub> per year may be emitted by fossil fuels expended for intensive confinement operations (Steinfeld et al. 2006). Energy uses in these industrial facilities differ substantially from those in smaller-scale, extensive or pasture-based farms. While a large portion goes toward heating, cooling, and ventilation systems, more than half of the energy used for intensive confinement operations is expended by feed crop production, specifically to produce seed, herbicides, and pesticides, as well as the fossil fuels used to operate farm machinery in the production of feed crops (Steinfeld et al. 2006).

According to the FAO's estimates, CO<sub>2</sub> emissions from farm animal processing total several tens of millions of tonnes per year (Steinfeld et al. 2006). The amount of fossil fuels burned varies depending on the species and type of animal product. For example, processing 1 kilogram of beef requires 4.37 megajoules, or 1.21 kilowatt-hours, while processing 1 dozen eggs requires more than 6 megajoules, or 1.66 kilowatt-hours (Steinfeld et al. 2006).

That same 1 kilogram of beef may result in GHGs equivalent to 36.4 kilograms of CO<sub>2</sub>, with almost all the energy consumed attributed to the production and transport of feed (Ogino et al. 2007).

Approximately 0.8 million tonnes of CO<sub>2</sub> are emitted annually from the transportation of feed and animal products to the places where they will be consumed (Steinfeld et al. 2006). Indeed, the international trade of meat alone can result in 500,000 to 850,000 tonnes of CO<sub>2</sub> annually (Steinfeld et al. 2006).

Farm animals and animal production facilities cover one-third of the planet's surface, using more than two-thirds of all available agricultural land (Haan et al. 1997). The livestock sector's CO<sub>2</sub> emissions from land use changes result from deforestation, land degradation, soil cultivation, and desertification.

A significant catalyst for the conversion of wooded areas to grazing land or cropland for feed production (Steinfeld et al. 2006), animal agriculture may emit 2.4 billion tonnes of CO<sub>2</sub> annually as a result of deforestation (Steinfeld et al. 2006). This sector has particularly devastated Latin America (Steinfeld et al. 2006), the region experiencing the largest net loss of forests and greatest releases of stored carbon into the atmosphere, resulting from disappearing vegetation (Steinfeld et al. 2006). One of the chief causes of Latin America's deforestation, according to the FAO, is cattle ranching (FAO 2005a).

Other important ecosystems are also threatened by increasing farm animal populations. Brazil's Cerrado region, the world's most biologically diverse savannah, produces half of the country's soy crops (Klink and Machado 2005; WWF 2007a, 2007b). As noted by the World Wildlife Fund (WWF 2007a), the region's animal species "are competing with the rapid expansion of Brazil's agricultural frontier, which focuses primarily on soy and corn. Ranching is another major threat to the region, as it produces almost 40 million cattle a year."

Farm animal production also results in releases of up to 28 million tonnes of CO<sub>2</sub> per year from cultivated soils (Steinfeld et al. 2006). Soils, like forests, act as carbon sinks and store more than twice the carbon found in vegetation or in the atmosphere (Steinfeld et al. 2006). Human activities, however, have significantly depleted the amount of carbon sequestered in the soil, contributing to GHG emissions.

Desertification, or the degradation of land in arid, semi-arid, and dry sub-humid areas, is also exacerbated and facilitated by the animal agriculture sector (FAO 2007). By reducing the productivity and amount of vegetative cover, desertification allows CO<sub>2</sub> to escape into the atmosphere.

Desertification of pastures due to animal agriculture is responsible for up to 100 million tonnes of CO<sub>2</sub> emissions annually (Steinfeld et al. 2006).

### ***Nitrogen from Fertilizer and Feed Production***

Feeding the global population of livestock requires at least 80 percent of the world's soybean crop and more than half of all corn (Ash M, Nierenberg D, personal communication; Halweil B, Smil V, personal communication), a plant whose growth is especially dependent on nitrogen-based artificial fertilizers. Natural sources of fixed nitrogen, the form easily available as fertilizer for plants, are limited, necessitating artificial fertilizer production. Prior to the development of the Haber-Bosch process, the amount of sustainable life on Earth was restricted by the amount of nitrogen made available to plants by bacteria and lightning. Modern fertilizer manufacturing, heavily reliant on fossil fuels, has taken a once-limited nutrient and made it available in massive quantities for crop farmers in the industrialized world and, increasingly, the developing world.

“The changes to the nitrogen cycle are larger in magnitude and more profound than the changes to the carbon cycle....But the nitrogen cycle is being neglected” (Bohan 2007), according to Elizabeth Holland, a Senior Scientist with the National Center for Atmospheric Research (NCAR). In addition, the Third International Nitrogen Conference’s co-chairs highlighted the role of farm animal production in the Nanjing Declaration on Nitrogen Management, a statement presented to the United Nations Environment Programme, recognizing that “a growing proportion of the world’s population consumes excess protein and calories, which may lead to human health problems. The associated production of these dietary proteins (especially animal products) leads to further disturbance of the nitrogen cycle” (Zhu et al. 2004).

According to University of Manitoba’s Vaclav Smil, a nitrogen cycle expert, “we have perturbed the global nitrogen cycle more than any other, even carbon” (Pollan 2006). Indeed, the overwhelming majority of all crops grown in the industrialized world are nitrogen-saturated, and overuse of nitrogen in crop production, nitrogen run-off into waterways, and the millions of tons of nitrogen found in farm animal manure threaten environmental integrity and public health.

### ***Methane and Nitrous Oxide***

The animal agriculture sector is also responsible for 35 to 40 percent of annual anthropogenic methane emissions (Steinfeld et al. 2006) that result from both enteric fermentation in ruminants and farm animal manure. Methane emissions are impacted by a number of factors including the animal’s age, bodyweight, feed quality, digestive efficiency, and exercise (Paustian et al. 2006; Steinfeld et al. 2006).

Ruminants emit methane as part of their digestive process, which involves microbial (enteric) fermentation (Steinfeld et al. 2006; U.S. EPA 2006). While individual animals produce relatively small

amounts of methane (U.S. EPA 2007b), the more than one billion ruminants, as documented by FAOSTAT, reared annually amount to a significant methane source. Indeed, enteric fermentation generates approximately 86 million tonnes of methane emissions worldwide (Steinfeld et al. 2006).

Typically, cattle confined in feedlots or in intensive confinement dairy operations are fed an unnatural diet of concentrated, high-protein feed comprised of corn and soybeans. While cattle may gain weight rapidly when fed this diet (Pollan 2002), it can cause a range of illnesses (Smith 1998). This diet may also lead to increased methane emissions. The standard diet fed to beef cattle confined in feedlots contributes to manure with a “high methane producing capacity” (U.S. EPA 1998). In contrast, cattle raised on pasture, eating a more natural, low-energy diet comprised of grasses and other forages, produce manure with about half of the potential to generate methane (U.S. EPA 1998).

Farm animals produce billions of tons of manure, with confined farm animals in the United States alone generating approximately 500 million tons of solid and liquid waste annually (U.S. EPA 2003). Storing and disposing of these immense quantities of manure can lead to significant anthropogenic emissions of methane and nitrous oxide (U.S. EPA 2007a). For example, according to the Pew Center on Global Climate Change, farm animal manure management accounts for 25 percent of agricultural methane emissions in the United States and 6 percent of agricultural nitrous oxide emissions (Paustian et al. 2006). Globally, emissions from pig manure alone account for almost half of all GHG emissions from farm animal manure (Steinfeld et al. 2006).

Farm animal manure is the source of almost 18 million tonnes of annual methane emissions (Steinfeld et al. 2006). Between 1990 and 2005 in the United States, methane emissions from dairy cow and pig manure rose by 50 and 37 percent, respectively (U.S. EPA 2007a). The U.S. EPA (2007a) traces this increase to the trend towards housing dairy cows and pigs in larger facilities that typically use liquid

manure management systems, which were first in use in the 1960s (Miner et al. 2000) but are now found in large dairy operations across the United States and in some developing countries, as well as in the majority of industrial pig operations worldwide.

While 70 percent of anthropogenic emissions of nitrous oxide result from crop and animal agriculture combined (Steinfeld et al. 2006), farm animal production accounts for 65 percent of global nitrous oxide emissions (Steinfeld et al. 2006). Manure and urine from ruminants, once deposited on the soil, emit nitrous oxide, and, in the United States, a 10-percent rise in nitrous oxide emissions between 1990 and 2005 can be traced, in part, to changes in the poultry industry, including an overall increase in the domestic stock of birds used for meat and egg production (U.S. EPA 2007a).

### ***Conflict, Hunger, and Disease***

As is the case with animal agriculture's impacts on soil, water, and air quality, the sector's contributions to climate change cannot be viewed in a vacuum. Climate change is having far-reaching consequences, perhaps most startlingly seen in growing conflicts among pastoral communities.

Environmental degradation has been cited as one of the catalysts for ongoing conflicts in Darfur and other areas of Sudan (UNEP 2007) where the effects of climate change have led to untenable conditions. As temperatures rise and water supplies dry up, farmers and herders are fighting to gain and control diminishing arable land and water (Baldauf 2006).

In a 2007 report issued by the United Nations Environment Programme, the agency tied two of its critical concerns in Sudan—land degradation and desertification—to “an explosive growth in livestock numbers.” In addition to citing climate change as one factor that led to the Darfur conflict (Ban 2007), United Nations Secretary-General Ban Ki-moon has noted that natural disasters, droughts, and other

changes brought about by global warming “are likely to become a major driver of war and conflict” (UN 2007).

According to the IPCC (2007a), many areas already suffering from drought will become drier, exacerbating the risks of both hunger and disease. By 2020, up to 250 million people may experience water shortages, and, in some countries, food production may be cut in half (IPCC 2007a). By 2050—the same year by which the FAO projects that meat and dairy production will double from present levels, primarily in the developing world (Steinfeld et al. 2006)—130 million people in Asia may suffer from climate change-related food shortages (Casey 2007).

Global temperature shifts may also hasten the speed at which infectious diseases emerge and re-emerge (Epstein and Mills 2005). According to the World Health Organization, “the chief risk factor for emerging zoonotic diseases is environmental degradation by humans, particularly deforestation, logging, and urbanisation” (Fleck 2004). The clear-cutting of forests for soybean cultivation, logging, and other industries enables viruses to exploit such newly exposed niches (Greger 2007).

### ***Strategies and Next Steps***

Mitigating the animal agriculture sector’s contributions to climate change necessitates comprehensive and immediate action by policy makers, producers, and consumers.

Enhanced regulation is required in order to hold facilities accountable for their GHG emissions. One critical step is accurately pricing environmental services, natural resources that are typically free or underpriced, leading to “overexploitation and pollution” (Steinfeld et al. 2006).

Thus far, most mitigation and prevention strategies undertaken by the animal agriculture sector have focused on technical solutions. For example, researchers are investigating the reformulation of ruminant diets to reduce enteric fermentation and some methane emissions (Connolly 2007). One such remedy is a plant-based bolus, formulated to reduce excessive fermentation and regulate the metabolic activity of rumen bacteria to reduce methane emissions from both the animals and their manure (Drochner W, Nierenberg, D, personal communication).

The USDA and U.S. EPA assist in funding anaerobic digester projects domestically and abroad (U.S. EPA 2007c; Sutherly 2007). These digesters, now in use at some large-scale intensive confinement facilities, capture methane from manure to use as a source of energy (Storck 2007), but are typically not economically viable for small-scale farms (Silverstein 2007).

In addition, producers are burning animal waste for fuel. The world's foremost pig producer, Smithfield Foods, and one of the top poultry producers, Tyson Foods, are both using animal byproduct fats to create biofuels (Johnston 2007; PR Newswire 2007).

McDonald's and agribusiness giant Cargill, which was supplying the company with soy for use as chicken feed, recently entered into an agreement with Brazil's other chief soy traders. Engineered by international environmental organization Greenpeace, a two-year moratorium is currently in place to prevent purchases of soy from the country's newly deforested areas (Kaufman 2007).

As consumers increasingly favor more environmentally friendly products and techniques, reducing consumption of meat, eggs, and milk, as well as choosing more sustainably produced animal products, such as those from organic systems, may prove equally critical strategies. Indeed, organic farming has the potential to reduce GHG emissions and sequester carbon, according to the International Federation

of Organic Agriculture Movements (IFOAM 2004), and raising cattle for beef organically on grass, in contrast to fattening confined cattle on concentrated feed, may emit 40-percent less GHGs and consume 85-percent less energy than conventionally produced beef (Cederberg and Stadig 2003; Fanelli 2007; Ogino et al. 2007).

There remains, however, an immediate need for more research regarding both technical and less technology-dependent strategies to record existing GHG emissions from individual production facilities and to provide lessons to producers and policy makers for reducing the climate damaging impacts of animal agriculture.

Given the urgency for global action, calls echoed by scientists and world leaders alike, individual consumers must also participate. The authors of a September 2007 *Lancet* article put forth several recommendations, including the reduction of meat and milk intake by high-income countries as “the urgent task of curtailing global greenhouse-gas emissions necessitates action on all major fronts,” and concluded that, for high-income countries, “greenhouse-gas emissions from meat-eating warrant the same scrutiny as do those from driving and flying” (McMichael et al. 2007).

## **Conclusion**

As the numbers of farm animals reared for meat, egg, and dairy production increase, so do their emissions. By 2050, global farm animal production is expected to double from present levels. The environmental impacts of animal agriculture require that governments, international organizations, producers, and consumers focus more attention on the role played by meat, egg, and dairy production. Mitigating and preventing the environmental harms caused by this sector require immediate and substantial changes in regulation, production practices, and consumption patterns.

## References

- Baldauf S. 2006. Africans are already facing climate change. *Christian Science Monitor* (Boston, MA) 6 November: 4.
- Ban K. 2007. A climate culprit in Darfur. *The Washington Post* (Washington, DC) 16 June: A15.
- Bierbaum RM, Holdren JP, MacCracken MC, Moss RH, Raven PH, eds. 2007. *Confronting Climate Change: Avoiding the Unmanageable, Managing the Unavoidable*. Report prepared for the United Nations Commission on Sustainable Development. Research Triangle Park, NC and Washington, DC:Sigma Xi and United Nations Foundation. Available: [http://www.unfoundation.org/files/pdf/2007/SEG\\_Report.pdf](http://www.unfoundation.org/files/pdf/2007/SEG_Report.pdf) [accessed 23 October 2007].
- Bohan S. 2007. Nitrogen overdose. *Oakland Tribune* (Oakland, CA) 12 August.
- Casey M. 2007. Report: millions face hunger from climate change. *The Christian Post* (Washington, DC) 10 April. Available: [http://www.christianpost.com/article/20070410/26802\\_Report:\\_Millions\\_Face\\_Hunger\\_from\\_Climate\\_Change.htm](http://www.christianpost.com/article/20070410/26802_Report:_Millions_Face_Hunger_from_Climate_Change.htm) [accessed 3 January 2008].
- Cederberg C, Stadig M. 2003. System expansion and allocation in life cycle assessment of milk and beef production. *Int J LCA* 8:350-356.
- Connolly K. 2007. Pill stops cow burps and helps save the planet. *The Guardian* (London, England) 23 March: 23.
- Epstein P, Mills E, eds. 2005. *Climate Change Futures: Health, Ecological, and Economic Dimensions*. Boston, MA:Center for Health and the Global Environment.
- Fanelli D. 2007. Meat is murder on the environment. *New Scientist*, 18 July: 15. Available: [http://environment.newscientist.com/article.ns?id=mg19526134.500&feedId=online-news\\_rss20](http://environment.newscientist.com/article.ns?id=mg19526134.500&feedId=online-news_rss20) [accessed 23 October 2007].
- Fleck F. 2004. Experts urge action to stop animal diseases infecting humans. *BMJ* 328:1158.

- FAO. 2007. Desertification. Available: <http://www.fao.org/desertification/default.asp?lang=en>  
[accessed 23 October 2007].
- FAO. 2005a. Cattle ranching is encroaching on forests in Latin America. Available:  
<http://www.fao.org/newsroom/en/news/2005/102924/index.html> [accessed 23 October 2007].
- FAO. 2005b. Responding to the “livestock revolution”—the case for livestock public policies.  
Available: [http://www.fao.org/ag/againfo/resources/documents/policies/briefs/01/EN/AGA01\\_10.pdf](http://www.fao.org/ag/againfo/resources/documents/policies/briefs/01/EN/AGA01_10.pdf) [accessed 23 October 2007].
- FAO. 2006. Livestock a major threat to the environment: remedies urgently needed. Available:  
<http://www.fao.org/newsroom/en/news/2006/1000448/index.html> [accessed 23 October 2007].
- Greger M. 2007. The human/animal interface: emergence and resurgence of zoonotic infectious diseases. *Crit Rev Microbiol* 33:243-299.
- Haan C, Steinfeld H, Blackburn H. 1997. *Livestock and the Environment: Finding a Balance*. Brussels, Belgium:European Commission Directorate-General for Development.
- IFOAM (International Federation of Organic Agriculture Movements). 2004. The role of organic agriculture in mitigating climate change. Available:  
[http://www.ifoam.org/press/positions/pdfs/Role\\_of\\_OA\\_mitigating\\_climate\\_change.pdf](http://www.ifoam.org/press/positions/pdfs/Role_of_OA_mitigating_climate_change.pdf)  
[accessed 23 October 2007].
- IPCC. 2007a. *Climate Change 2007: Climate Change Impacts, Adaptation and Vulnerability, Summary for Policymakers, Working Group II Contribution to the Intergovernmental Panel on Climate Change Fourth Assessment Report*. Brussels, Belgium: Intergovernmental Panel on Climate Change.
- IPCC. 2007b. *Climate Change 2007: Summary for Policymakers of the Synthesis Report of the IPCC Fourth Assessment Report*. Draft copy, 16 November.

- IPCC. 2007c. Climate Change 2007: The Physical Science Basis, Summary for Policymakers, Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Paris, France: Intergovernmental Panel on Climate Change.
- Johnston T. 2007. Tyson teams with ConocoPhillips to produce renewable diesel fuel. MeatingPlace.com 16 April. Available: <http://meatingplace.com/MembersOnly/webNews/details.aspx?item=17806> [accessed 23 October 2007].
- Kaufman M. 2007. New allies on the Amazon: McDonald's, Greenpeace unite to prevent rainforest clearing. The Washington Post (Washington, DC) 24 April: D01.
- Klink CA, Machado RB. 2005. Conservation of the Brazilian Cerrado. *Conserv Biol* 19:707-713.
- McMichael AJ, Powles JW, Butler CD, Uauy R. 2007. Food, livestock production, energy, climate change, and health. *Lancet* 370:1253–1263; doi:10.1016/S0140-6736(07)61256-2 [Online 13 September 2007].
- Miner J, Humenik F, Overcash R. 2000. Managing Livestock Wastes to Preserve Environmental Quality. Ames, Iowa: Iowa University Press.
- NASA (National Aeronautics and Space Administration). 2006. 2005 warmest year in over a century. Available: [http://www.nasa.gov/vision/earth/environment/2005\\_warmest.html](http://www.nasa.gov/vision/earth/environment/2005_warmest.html) [accessed 23 October 2007].
- Nierenberg D. 2006. Rethinking the global meat industry. In: State of the World 2006: A Worldwatch Institute Report on Progress Toward a Sustainable Society (Starke L, ed). New York: W.W. Norton & Company, Inc., 24-40.
- Ogino A, Orito H, Shimada K, Hirooka H. 2007. Evaluating environmental impacts of the Japanese beef cow-calf system by the life cycle assessment method. *Animal Sci J* 78:424-432.
- Paustian K, Antle M, Sheehan J, Eldor P. 2006. Agriculture's Role in Greenhouse Gas Mitigation. Washington, DC: Pew Center on Global Climate Change.

- Pollan M. 2002. Power steer. *The New York Times Magazine*, 31 March: 44.
- Pollan M. 2006. *The Omnivore's Dilemma: A Natural History of Four Meals*. New York: The Penguin Press.
- PR Newswire. 2007. Smithfield joins the Chicago Climate Exchange. Available: <http://sev.prnewswire.com/food-beverages/20070226/CGM04426022007-1.html> [accessed 24 October 2007].
- Silverstein K. 2007. The appeal of animal waste. *EnergyBiz Insider*, 10 August. Available: [http://www.energycentral.com/centers/energybiz/ebi\\_detail.cfm?id=367](http://www.energycentral.com/centers/energybiz/ebi_detail.cfm?id=367) [accessed 8 October 2007].
- Smith RA. 1998. Impact of disease on feedlot performance: a review. *J Anim Sci* 76:272-274.
- Steinfeld H, Gerber P, Wassenaar T, Castel V, Rosales M, de Haan C. 2006. *Livestock's Long Shadow: Environmental Issues and Options*. Rome, Italy: Food and Agriculture Organization of the United Nations.
- Storck AB. 2007. More farms find unlikely power source: manure. *Meatingplace.com* 30 July. Available: <http://www.meatingplace.com/MembersOnly/webNews/details.aspx?item=18539> [accessed 24 October 2007].
- Sutherly B. 2007. Ohio farms planning to use cows, chickens to generate energy. *Dayton Daily News* (Dayton, OH), 22 June. Available: <http://www.daytondailynews.com/n/content/oh/story/news/local/2007/07/21/ddn072207farmenergy.html> [accessed 3 January 2008].
- Thorne PS. 2007. Environmental health impacts of concentrated animal feeding operations: anticipating hazards – searching for solutions. *Environ Health Perspect* 115:296-297.
- Topping Jr. JC. 2007. Summit aftermath: study by NASA and university scientists shows world temperature reaching a level not seen in thousands of years and raises grave concern of irreparable harm. Available:

[http://www.climate.org/programs/washington\\_summit\\_temperature\\_rise.shtml](http://www.climate.org/programs/washington_summit_temperature_rise.shtml) [accessed 24 October 2007].

UN (United Nations). 2007. Ban Ki-moon calls on new generation to take better care of Planet Earth than his own. Available:

<http://www.un.org/apps/news/story.asp?NewsID=21720&Cr=global&Cr1=warming> [accessed 24 October 2007].

UNEP (United Nations Environment Programme). 2007. Sudan: post-conflict environmental assessment. Available: [http://sudanreport.unep.ch/UNEP\\_Sudan.pdf](http://sudanreport.unep.ch/UNEP_Sudan.pdf) [accessed 24 October 2007].

USDA. 2004. U.S. Agriculture and Forestry Greenhouse Gas Inventory: 1990-2001. Washington, DC:U.S. Department of Agriculture.

U.S. EPA. 1998. Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990-1996. Washington, DC:U.S. Environmental Protection Agency.

U.S. EPA. 2003. National Pollutant Discharge Elimination System Permit Regulation and Effluent Limitation Guidelines and Standards for Concentrated Animal Feeding Operations (CAFOs); Final Rule. Fed Reg 68:7175-7274.

U.S. EPA. 2006. Methane: sources and emissions. Available: <http://www.epa.gov/methane/sources.html> [accessed 24 October 2007].

U.S. EPA. 2007a. Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990-2005. Washington, DC:U.S. Environmental Protection Agency.

U.S. EPA. 2007b. Ruminant livestock: frequent questions. Available: <http://www.epa.gov/methane/rlep/faq.html> [accessed 24 October 2007].

U.S. EPA. 2007c. U.S. government accomplishments in support of the Methane to Markets partnership. Available: <http://www.epa.gov/methanetomarkets/accompreport.htm> [accessed 4 October 2007].

Verge XPC, De Kimpe C, Desjardins RL. 2007. Agricultural production, greenhouse gas emissions and mitigation potential. *Agr Forest Meteorol* 142:255-269.

WWF (World Wildlife Fund). 2007a. Brazilian savannas. Available:

[http://www.panda.org/news\\_facts/education/best\\_place\\_species/current\\_top\\_10/brazilian\\_savannas\\_.cfm](http://www.panda.org/news_facts/education/best_place_species/current_top_10/brazilian_savannas_.cfm) [accessed 24 October 2007].

WWF (World Wildlife Fund). 2007b. Facts about soy production and the Basel Criteria. Available:

[http://assets.panda.org/downloads/factsheet\\_soy\\_eng.pdf](http://assets.panda.org/downloads/factsheet_soy_eng.pdf) [accessed 24 October 2007].

Zhu Z, Minami K, Galloway J. 2004. Nanjing Declaration on nitrogen management, presented to the United Nations Environment Programme. Available:

[http://www.initrogen.org/fileadmin/user\\_upload/nanjing/nanjing\\_declaration-041016.pdf](http://www.initrogen.org/fileadmin/user_upload/nanjing/nanjing_declaration-041016.pdf) [accessed 24 October 2007].

